

**Allergy and Anaphylaxis
Medication Administration Authorization Plan**

Place Child's Picture
Here (optional)

This form must be completed fully in order for Child Care Providers/staff to administer the required medication and follow the plan. **This authorization is NOT TO EXCEED 1 YEAR.**
Page 1 to be completed by the Authorized Health Care Provider.
FOR ALLERGY AND ANAPHYLAXIS MEDICATION ONLY - THIS FORM REPLACES OCC 1216

CHILD'S NAME: _____ Date of Birth: ____/____/____ **Date of plan:** _____
 Child has **Allergy** to _____ Ingestion/Mouth Inhalation Skin Contact Sting Other _____
 Child has had anaphylaxis: Yes No
 Child has asthma: Yes No (If yes, higher chance severe reaction) Child
 may self-carry medication: Yes No
 Child may self-administer medication: Yes No

Allergy and Anaphylaxis Symptoms	Treatment Order	
If child has ingested a food allergen, been stung by a bee or exposed to an allergy trigger	Antihistamine :Oral /By Mouth <input type="checkbox"/> Call Parent <input type="checkbox"/> Call 911	Epinephrine(EpiPen) IM Injection in Thigh <input type="checkbox"/> Call 911 <input type="checkbox"/> Call Parent
is Not exhibiting or complaining of any symptoms, OR		
Exhibits or complains of any symptoms below:		
Mouth: itching, tingling, swelling of lips, tongue ("mouth feels funny")		
Skin: hives, itchy rash, swelling of the face or extremities		
Throat*: difficulty swallowing ("choking feeling"), hoarseness, hacking cough		
Lung*: shortness of breath, repetitive coughing, wheezing		
Heart*: weak or fast pulse, low blood pressure, fainting, pale, blueness		
Gut: nausea, abdominal cramps, vomiting, diarrhea		
Other:		
If reaction is progressing (several of the above areas affected)		

Potentially life threatening. The severity of symptoms can quickly change

Medication	Medication: Brand and Strength	Dose	Route	Frequency
Epinephrine(EpiPen)				
Antihistamine				
Other:				

EMERGENCY Response:

- 1) **Inject epinephrine right away! Note time when epinephrine was administered.**
- 2) Call 911: Ask for ambulance with epinephrine. Advise rescue squad when epinephrine was given. Stay with child.
- 3) Call parents. Advise parent of the time that epinephrine was given and 911 was called.
- 4) Keep child lying on his/her back. If the child vomits or has trouble breathing, place child on his/her side.
- 5) Give other medicine, if prescribed.

PRESCRIBER'S NAME/TITLE		Place stamp here
TELEPHONE	FAX	
ADDRESS		
PRESCRIBER'S SIGNATURE (Parent/guardian cannot sign here) (original signature or signature stamp only)		DATE (mm/dd/yyyy)

Maryland State Department of Education
Office of Child Care
Allergy and Anaphylaxis
Medication Administration Authorization Plan

Child's Name: _____ Date of Birth: _____

PARENT/GUARDIAN AUTHORIZATION			
I request the authorized child care staff to administer the medication or to supervise the child in self-administration as prescribed above. I certify that I have legal authority to consent to medical treatment for the child named above, including the administration of medication at the facility. I understand that at the end of the authorized period an authorized individual must pick up the medication; otherwise, it will be discarded. I authorize child care staff and the authorized prescriber indicated on this form to communicate in compliance with HIPAA. I understand that per COMAR 13A.15, 13A.16, 13A.17, and 13A.18, the child care program may revoke the child's authorization to self-carry/self-administer medication.			
PARENT/GUARDIAN SIGNATURE		DATE (mm/dd/yyyy)	INDIVIDUALS AUTHORIZED TO PICK UP MEDICATION
CELL PHONE #		HOME PHONE #	WORK PHONE #
Emergency Contact(s)	Name/Relationship	Phone Number to be used in case of Emergency	
Parent/Guardian 1			
Parent/Guardian 2			
Emergency 1			
Emergency 2			
Section IV. CHILD CARE STAFF USE ONLY			
Child Care Responsibilities:	1. Medication named above was received	<input type="checkbox"/> Yes	<input type="checkbox"/> No
	2. Medication labeled as required by COMAR	<input type="checkbox"/> Yes	<input type="checkbox"/> No
	3. OCC 1214 Emergency Card updated	<input type="checkbox"/> Yes	<input type="checkbox"/> No
	4. OCC 1215 Health Inventory updated	<input type="checkbox"/> Yes	<input type="checkbox"/> No
	5. Modified Diet/Exercise Plan	<input type="checkbox"/> Yes	<input type="checkbox"/> No <input type="checkbox"/> N/A
	6. Individualized Plan: IEP/IFSP	<input type="checkbox"/> Yes	<input type="checkbox"/> No <input type="checkbox"/> N/A
	7. Staff approved to administer medication is available onsite, field trips	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Reviewed by (printed name and signature):			DATE (mm/dd/yyyy)

DOCUMENT MEDICATION ADMINISTRATION HERE

DATE	TIME	MEDICATION	DOSAGE	ROUTE	REACTIONS OBSERVED (IF ANY)	SIGNATURE



FARE
Food Allergy Research & Education

FOOD ALLERGY & ANAPHYLAXIS EMERGENCY CARE PLAN

**PLACE
PICTURE
HERE**

Name: _____ D.O.B.: _____

Allergic to: _____

Weight: _____ lbs. Asthma: **Yes (higher risk for a severe reaction)** **No**

NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE.

Special Situation/Circumstance - If this box is checked, the child has an extremely severe allergy to the following food(s) _____.
Even if the child has MILD symptoms after eating (ingesting) this food(s), Give Epinephrine immediately.

For ANY of the following SEVERE SYMPTOMS



LUNG

Shortness of breath, wheezing, repetitive cough



HEART

Pale or bluish skin, faintness, weak pulse, dizziness



THROAT

Tight or hoarse throat, trouble breathing or swallowing



MOUTH

Significant swelling of the tongue or lips



SKIN

Many hives over body, widespread redness



GUT

Repetitive vomiting, severe diarrhea



OTHER

Feeling something bad is about to happen, anxiety, confusion

OR A COMBINATION

of symptoms from different body areas



- 1. INJECT EPINEPHRINE IMMEDIATELY.**
- 2. Call 911.** Tell emergency dispatcher the person is having anaphylaxis and may need epinephrine when emergency responders arrive.
 - Consider giving additional medications following epinephrine:
 - » Antihistamine
 - » Inhaler (bronchodilator) if wheezing
 - Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
 - If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
 - Alert emergency contacts.
 - Transport patient to ER, even if symptoms resolve. Patient should remain in ER for at least 4 hours because symptoms may return

MILD SYMPTOMS



NOSE

Itchy or runny nose, sneezing



MOUTH

Itchy mouth



SKIN

A few hives, mild itch



GUT

Mild nausea or discomfort

FOR MILD SYMPTOMS FROM MORE THAN ONE BODY SYSTEM, GIVE EPINEPHRINE.

FOR MILD SYMPTOMS FROM A SINGLE BODY SYSTEM (E.G. SKIN, GI, ETC.), FOLLOW THE DIRECTIONS BELOW:

1. Antihistamines may be given, if ordered by a healthcare provider.
2. Stay with the person; alert emergency contacts.
3. Watch closely for changes. If symptoms worsen, give epinephrine.

MEDICATIONS/DOSES

Epinephrine Brand or Generic: _____

Epinephrine Dose: 0.1 mg IM 0.15 mg IM 0.3 mg IM

Antihistamine Brand or Generic: _____

Antihistamine Dose: _____

Other (e.g., inhaler-bronchodilator if wheezing): _____

PATIENT OR PARENT/GUARDIAN AUTHORIZATION SIGNATURE

DATE


HEALTHCARE PROVIDER AUTHORIZATION SIGNATURE

DATE




HOW TO USE AUVI-Q® (EPINEPHRINE INJECTION, USP), KALEO

1. Remove Auvi-Q® from the outer case. Pull off red safety guard.
2. Place black end of Auvi-Q® against the middle of the outer thigh.
3. Press firmly until you hear a click and hiss sound, and hold in place for 2 seconds.
4. Call 911 and get emergency medical help right away.




HOW TO USE EPIPEN®, EPIPEN JR® (EPINEPHRINE) AUTO-INJECTOR AND EPINEPHRINE INJECTION

1. (AUTHORIZED GENERIC OF EPIPEN®), USP AUTO-INJECTOR, MYLAN AUTO-INJECTOR, MYLAN
2. Remove the EpiPen® or EpiPen Jr® Auto-Injector from the clear carrier tube.
3. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, remove the blue safety release by pulling straight up.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
5. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.




HOW TO USE IMPAX EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF ADRENACLICK®), USP AUTO-INJECTOR, AMNEAL PHARMACEUTICALS

1. Remove epinephrine auto-injector from its protective carrying case.
2. Pull off both blue end caps: you will now see a red tip. Grasp the auto-injector in your fist with the red tip pointing downward.
3. Put the red tip against the middle of the outer thigh at a 90-degree angle, perpendicular to the thigh. Press down hard and hold firmly against the thigh for approximately 10 seconds.
4. Remove and massage the area for 10 seconds. Call 911 and get emergency medical help right away.



HOW TO USE TEVA'S GENERIC EPIPEN® (EPINEPHRINE INJECTION, USP) AUTO-INJECTOR, TEVA PHARMACEUTICAL INDUSTRIES

1. Quickly twist the yellow or green cap off of the auto-injector in the direction of the "twist arrow" to remove it.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, pull off the blue safety release.
3. Place the orange tip against the middle of the outer thigh at a right angle to the thigh.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
5. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.



HOW TO USE SYMJEPI™ (EPINEPHRINE INJECTION, USP)

1. When ready to inject, pull off cap to expose needle. Do not put finger on top of the device.
2. Hold SYMJEPI™ by finger grips only and slowly insert the needle into the thigh. SYMJEPI™ can be injected through clothing if necessary.
3. After needle is in thigh, push the plunger all the way down until it clicks and hold for 2 seconds.
4. Remove the syringe and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.
5. Once the injection has been administered, using one hand with fingers behind the needle slide safety guard over needle.



ADMINISTRATION AND SAFETY INFORMATION FOR ALL AUTO-INJECTORS:

1. Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than mid-outer thigh. In case of accidental injection, go immediately to the nearest emergency room.
2. If administering to a young child, hold their leg firmly in place before and during injection to prevent injuries.
3. Epinephrine can be injected through clothing if needed.
4. Call 911 immediately after injection.

OTHER DIRECTIONS/INFORMATION (may self-carry epinephrine, may self-administer epinephrine, etc.):

Epinephrine first, then call 911. Monitor the patient and call their emergency contacts right away.

EMERGENCY CONTACTS – CALL 911

RESCUE SQUAD: _____
DOCTOR: _____ PHONE: _____
PARENT/GUARDIAN: _____ PHONE: _____

OTHER EMERGENCY CONTACTS

NAME/RELATIONSHIP: _____ PHONE: _____
NAME/RELATIONSHIP: _____ PHONE: _____
NAME/RELATIONSHIP: _____ PHONE: _____